



SCIENCE @ DIRECT

Register or Login: Password: Athens Login[Home](#)[Journals](#)[Abstract Databases](#)[Books](#)[Reference Works](#)[My Profile](#)[Alerts](#)[? Help](#)WELCOME GUEST USER [info](#)**Tetrahedron Letters**

Volume 31, Issue 11, 1990, Pages 1517-1520

doi:10.1016/0040-4039(90)80004-6 [? Cite or link using doi](#)
Copyright © 1990 Published by Elsevier Science Ltd. All rights reserved.

This Document

- ▶ **Abstract**
- [Abstract + References](#)
- [PDF \(262 K\)](#)

Comparison of three methods for the synthesis of carborane carboxylic acid esters

Stephen B. Kahl

Department of Pharmaceutical Chemistry University of California, San Francisco San Francisco, CA 94143-0446, USA

Received 18 December 1989. Available online 15 March 2001.

Actions

- [E-mail Article](#)

Abstract

Three procedures for the esterification of polyhedral carborane carboxylic acids with long chain unsaturated fatty alcohols are compared with regard to rate of reaction, ease of isolation and over-all yield. The optimum procedure is based on room temperature reaction of the acid chloride and alcohol in the presence of 4-dimethylaminopyridine in CH_2Cl_2 .

Graphical Abstract

Esterification of 1,2-dicarba-closo-dodecaboranyl monocarboxylic acid with ten unsaturated fatty alcohols, as exemplified here with palmitoleyl alcohol, occurs most efficiently *via* reaction of the acid chloride and alcohol in the presence of p-dimethylamino pyridine.

**Tetrahedron Letters**

Volume 31, Issue 11, 1990, Pages 1517-1520

This Document

- ▶ **Abstract**
- [Abstract + References](#)
- [PDF \(262 K\)](#)

Actions

- [E-mail Article](#)

Home	Journals	Abstract Databases	Books	Reference Works	My Profile	Alerts	? Help
----------------------	--------------------------	------------------------------------	-----------------------	---------------------------------	----------------------------	------------------------	------------------------

Send [feedback](#) to ScienceDirect

Software and compilation © 2004 ScienceDirect. All rights reserved.

ScienceDirect® is a registered trademark of Elsevier B.V.

Your use of this service is governed by [Terms and Conditions](#). Please review our [Privacy Policy](#) for details on how we protect information that you supply.



SCIENCE @ DIRECT

Register or Login: Password: Athens L

[Home](#) [Journals](#) [Abstract Databases](#) [Books](#) [Reference Works](#) [My Profile](#) [Alerts](#) [Help](#)

WELCOME GUEST USER [info](#)

Biochemical and Biophysical Research Communications

Volume 262, Issue 1, 19 August 1999, Pages 275-284

doi:10.1006/bbrc.1999.1105 [Cite or link using doi](#)
Copyright © 1999 Academic Press. All rights reserved.

This Document

- ▶ **Abstract**
- [Abstract + References](#)
- [PDF \(126 K\)](#)

Actions

[E-mail Article](#)

Regular Article

Unsaturated Long-Chain N-Acyl-vanillyl-amides (N-AVAMs): Vanilloid Receptor Ligands That Inhibit Anandamide-Facilitated Transport and Bind to CB1 Cannabinoid Receptors^{*1}

Dominique Melck^{a, 1}, Tiziana Bisogno^{a, 1}, Luciano De Petrocellis^b, Huai-hu Chuang^c, David Julius^c, Maurizio Bifulco^d and Vincenzo Di Marzo^{a, 2}

Istituto per la Chimica di Molecole di Interesse Biologico³

^b Istituto di Cibernetica, Consiglio Nazionale delle Ricerche, Via Toiano 6, 80072, Arco Felice, Napoli, Italy

^c Department of Cellular and Molecular Pharmacology, University of California, San Francisco, California, 94143-0450

^d C.E.O.S., CNR, and Dipartimento di Biologia e Patologia Cellulare e Molecolare, Università di Napoli Federico II, 80131, Napoli, Italy

Received 29 June 1999. Available online 2 April 2002.

Abstract

We investigated the effect of changing the length and degree of unsaturation of the fatty acyl chain of *N*-(3-methoxy-4-hydroxy)-benzyl-*cis*-9-octadecenoamide (olvanil), a ligand of vanilloid receptors, on its capability to: (i) inhibit anandamide-facilitated transport into cells and enzymatic hydrolysis, (ii) bind to CB1 and CB2 cannabinoid receptors, and (iii) activate the VR1 vanilloid receptor. Potent inhibition of [¹⁴C]anandamide accumulation into cells was achieved with C20:4 n-6, C18:3 n-6 and n-3, and C18:2 n-6 *N*-acyl-vanillyl-amides (N-AVAMs). The saturated analogues and Δ^9 -*trans*-olvanil were inactive. Activity in CB1 binding assays increased when increasing the number of *cis*-double bonds in a n-6 fatty acyl chain and, in saturated N-AVAMs, was not greatly sensitive to decreasing the

chain length. The C20:4 n-6 analogue (arvanil) was a potent inhibitor of anandamide accumulation ($IC_{50} = 3.6 \mu M$) and was 4-fold more potent than anandamide on CB1 receptors ($K_i = 0.25-0.52 \mu M$), whereas the C18:3 n-3 N-AVAM was more selective than arvanil for the uptake ($IC_{50} = 8.0 \mu M$) vs CB1 receptors ($K_i = 3.4 \mu M$). None of the compounds efficiently inhibited [^{14}C]anandamide hydrolysis or bound to CB2 receptors. All N-AVAMs activated the cation currents coupled to VR1 receptors overexpressed in *Xenopus* oocytes. In a simple, intact cell model of both vanilloid- and anandamide-like activity, i.e., the inhibition of human breast cancer cell (HBCC) proliferation, arvanil was shown to behave as a "hybrid" activator of cannabinoid and vanilloid receptors.

Author Keywords: cannabinoid; endocannabinoid; capsaicin; analgesics; breast cancer cells; carrier

*1 Abbreviations used: N-AVAM, *N*-acyl-vanillyl-amide; VR1, vanilloid receptor type 1; RBL, rat basophilic leukemia; CB1, CB2, cannabinoid receptor type 1 and 2; TRP, transient receptor potential; TRPL, transient receptor potential-like; RTX, resiniferatoxin; FAAH, fatty acid amide hydroxylase; AM404, *N*-(4-hydroxyphenyl)-arachidonylamide; anandamide, *N*-arachidonoyl-ethanolamine; olvanil, *N*-(3-methoxy-4-hydroxy)-benzyl-*cis*-9-octadecenoamide; palvanil, *N*-(3-methoxy-4-hydroxy)-benzyl-hexadecanamide; arvanil, *N*-(3-methoxy-4-hydroxy)-benzyl-arachidonylamide; pseudocapsaicin, *N*-(3-methoxy-4-hydroxy)-benzyl-nonanamide; HBCC, human breast cancer cell

¹ These authors contributed equally to this work.

² Address correspondence and reprint requests to this author. Fax: +39-081-8041770. E-mail: VDM@TRINC.ICMIB.NA.CNR.IT.

³ Affiliated with the National Institute for the Chemistry of Biological Systems, Consiglio Nazionale delle Ricerche.

Biochemical and Biophysical Research Communications
Volume 262, Issue 1, 19 August 1999, Pages 275-284

This Document

• **Abstract**

• [Abstract + References](#)

• [PDF \(126 K\)](#)

Actions

• [E-mail Article](#)

[Home](#)

[Journals](#)

[Abstract Databases](#)

[Books](#)

[Reference Works](#)

[My Profile](#)

[Alerts](#)

[? Help](#)

[Send feedback](#) to ScienceDirect

Software and compilation © 2004 ScienceDirect. All rights reserved.

ScienceDirect® is a registered trademark of Elsevier B.V.

Your use of this service is governed by [Terms and Conditions](#). Please review our [Privacy Policy](#) for details on how we protect information that you supply.